

THREE SPHERES THEOREM FOR p -HARMONIC FUNCTIONS

VLADIMIR M. MIKLYUKOV, ANTTI RASILA AND MATTI VUORINEN

ABSTRACT. Three spheres theorem type result is proved for the p -harmonic functions defined on the complement of k -balls in the Euclidean n -dimensional space.

1. INTRODUCTION

A classical theorem by J. Hadamard gives the following relation between the maximum absolute values of an analytic function on three concentric circles.

1.1. Theorem. *Let $R_1 < r_1 < r_2 < r_3 < R_2$ and let f be an analytic function in the annulus $\{z \in \mathbb{C} : R_1 < |z| < R_2\}$. Denote the maximum of $|f(z)|$ on the circle $|z| = r$ by $M(r)$. Then*

$$M(r_2)^{\log(r_3/r_1)} \leq M(r_1)^{\log(r_3/r_2)} M(r_3)^{\log(r_2/r_1)}.$$

This result, known as the three circles theorem, was given by Hadamard without proof in 1896 [3]. For a discussion of the history of this result, see e.g. [8] and [5, pp. 323–325]. It is a natural question, what results of this type can be proved for other classes of functions. For example, a version of Hadamard's theorem can be proved for subharmonic functions in $\mathbb{R}^n$, $n \geq 2$, see [7, pp. 128–131].

Some generalizations of the three circles theorem will be studied here. For the formulation of our main result, Theorem 2.1, we recall some standard notation and definitions from the book [4]. We will consider solutions $v: \Omega \rightarrow \mathbb{R}$ of the p -Laplace equation

$$(1.2) \quad \operatorname{div}(|\nabla v|^{p-2} \nabla v) = 0, \quad 1 < p < \infty,$$

on an open set $\Omega \subset \mathbb{R}^n$ in the sense that will be described shortly. When $p = 2$ equation (1.2) reduces to the Laplace equation $\Delta u = 0$, whose solutions, harmonic functions, are studied in the classical potential theory. When $p \neq 2$ equation (1.2) is nonlinear and degenerates at the zeros of the gradient of v . It follows that the solutions, p -harmonic functions, need not be in $C^2(\Omega)$ and the equation must be understood in the weak sense. A weak solution of (1.2) is a function v in the Sobolev space $W_{\text{loc}}^{1,p}(\Omega)$ such that

$$(1.3) \quad \int_{\Omega} \langle |\nabla v|^{p-2} \nabla v, \nabla \varphi \rangle \, dm = 0$$

for all $\varphi \in C_0^\infty(\Omega)$, where $\langle \cdot, \cdot \rangle$ denotes the scalar product of vectors in $\mathbb{R}^n$, and m is the Lebesgue measure in $\mathbb{R}^n$.

1991 *Mathematics Subject Classification.* 35J60, 35B05, 35B50.

Key words and phrases. three circles theorem, p -harmonic functions, p -Laplacian.

The research was supported by the Academy of Finland, grant 107317 and by the Foundation Vilho, Yrjö ja Kalle Väisälän rahasto of Finnish Academy of Science and Letters.




FIGURE 1. 1-annulus $D_{\alpha,\beta}$ in $\mathbb{R}^2$ (left) and 2-annulus $D_{\alpha,\beta}$ in $\mathbb{R}^3$ (right).

It is easy to see that for all $\varphi \in C_0^\infty(\Omega)$ and $v \in C^2(\Omega)$,

$$\int_{\Omega} \langle |\nabla v|^{p-2} \nabla v, \nabla \varphi \rangle dm = - \int_{\Omega} \varphi \operatorname{div} (|\nabla v|^{p-2} \nabla v) dm$$

and, consequently, each C^2 -solution to (1.2) is a weak solution to (1.2).

Fix an integer k , $1 \leq k \leq n$ and a real number $t \geq 0$. The sets $B_k(t) = \{x \in \mathbb{R}^n : d_k(x) < t\}$ and $\Sigma_k(t) = \{x \in \mathbb{R}^n : d_k(x) = t\} = \partial B_k(t)$, where $d_k(x) = \left(\sum_{i=1}^k x_i^2\right)^{1/2}$, are respectively called k -ball and k -sphere in $\mathbb{R}^n$. For $k = n$ the k -ball $B_k(t)$ coincides with the standard Euclidean ball $B^n(t)$ and the k -sphere $\Sigma_k(t)$ is the Euclidean sphere $S^{n-1}(t)$. In particular, the symbol $\Sigma_k(0)$ below denotes the k -sphere with the radius 0, i.e.

$$\Sigma_k(0) = \{x = (x_1, \dots, x_k, \dots, x_n) : x_1 = \dots = x_k = 0\}.$$

Let $0 < \alpha < \beta < \infty$ be fixed and let

$$D_{\alpha,\beta} = \{x \in \mathbb{R}^n : \alpha < d_k(x) < \beta\}.$$

For $k = 1$ the set $D_{\alpha,\beta}$ is the union of the two layers between two parallel hyperplanes. For $1 < k < n$ the boundary of the domain $D_{\alpha,\beta}$ consists of two coaxial cylindrical surfaces.

Let $v \in C^0(D_{r,R})$, and let $M(r) = \limsup_{z \rightarrow \Sigma_k(r)} v(z)$. Suppose that $M(R) > M(r)$. Consider the function

$$v_{r,R}(x) = \frac{v(x) - M(r)}{M(R) - M(r)},$$

for $r < R$. Clearly, $\limsup_{z \rightarrow \Sigma_k(r)} v_{r,R}(z) \leq 0$ and $\limsup_{z \rightarrow \Sigma_k(R)} v_{r,R}(z) \leq 1$. Let

$$\xi(r, t) = \int_r^t s^{(1-k)/(p-1)} ds, \text{ and } u_0^{k,p}(t) = \frac{\xi(r, t)}{\xi(r, R)}.$$

Let $u(x) = u_0^{k,p}(d_k(x))$ for $x \in D_{r,R}$. It is clear (see Lemma 3.5) that u is a C^2 -solution to (1.2). We have

$$u(x)|_{\Sigma_k(r)} \equiv 0, \quad u(x)|_{\Sigma_k(R)} \equiv 1,$$

and

$$(1.4) \quad u(x) \geq v_{r,R}(x) \text{ if } x \in \Sigma_k(r) \text{ or } x \in \Sigma_k(R).$$

2. MAIN RESULTS

We will prove the following Hadamard type theorem for the p -harmonic functions defined on the complement of a k -ball. We use the method of proof from [6].

2.1. Theorem. *Let $1 < p < \infty$, $R > r > 0$ and let $v(x) \in W_{\text{loc}}^{1,p}(D_{r,\infty})$ be a continuous weak solution of (1.2) such that*

$$(2.2) \quad \int_r^\infty dt \left(\int_{\Sigma_k(t)} |v_{r,R} - u|^2 (|\nabla v_{r,R}|^{p-2} + |\nabla u|^{p-2}) d\mathcal{H}^{n-1} \right)^{-1} = \infty,$$

where $\mathcal{H}^{n-1}$ is the $(n-1)$ -dimensional Hausdorff measure. Then for all $t \in (r, R)$,

$$(2.3) \quad M(t) \leq (M(R) - M(r))u_0^{k,p}(t) + M(r).$$

Note that for $k = n$ (2.3) follows immediately from the comparison principle, see [4, p. 133].

2.4. Corollary. *Let $1 < p < \infty$, $R > r > 0$ and let $v(x) \in W_{\text{loc}}^{1,p}(D_{r,\infty})$, be a continuous weak solution of (1.2) such that*

$$(2.5) \quad \lim_{S \rightarrow \infty} \frac{1}{S^2} \int_{D_{r,S}} |v_{r,R} - u|^2 (|\nabla v_{r,R}|^{p-2} + |\nabla u|^{p-2}) dm = 0.$$

Then for all $t \in (r, R)$ the inequality (2.3) holds.

2.6. Corollary. *Let $1 < p < \infty$, $R > r > 0$ and let $v(x) \in W_{\text{loc}}^{1,p}(D_{r,\infty})$ be a continuous weak solution of (1.2) such that*

$$\int_{D_{r,\infty}} |v_{r,R} - u|^2 (|\nabla v_{r,R}|^{p-2} + |\nabla u|^{p-2}) dm \leq M < \infty.$$

Then for all $t \in (r, R)$ the inequality (2.3) holds.

For the formulation of a result of S. Granlund [2], Theorem 2.7 below, we introduce some notation and terminology. Let $p > 1$, $\Omega \subset \mathbb{R}^n$ be a bounded domain and let $F: \Omega \times \mathbb{R}^n \rightarrow \mathbb{R}$ be such that the following conditions hold.

(1) There are constants $\beta > \alpha > 0$ such that for a.e. $x \in \Omega$

$$\alpha|z|^p \leq F(x, z) \leq \beta|z|^p.$$

(2) For a.e. $x \in \Omega$ the function $z \mapsto F(x, z)$ is convex.

(3) The function $x \mapsto F(x, \nabla u(x))$ is measurable for all $u \in W^{1,p}(\Omega)$.

Let

$$I(u) = \int_{\Omega} F(x, \nabla u(x)) dm.$$

A function $u \in W^{1,p}(\Omega)$ is a *subminimum* in Ω if $I(u) \leq I(u - \eta)$ for all non-negative $\eta \in W_0^{1,p}(\Omega)$. Let

$$M(r) = \text{ess sup}_{x \in \overline{B}^n(r)} u(x), \quad \overline{B}^n(r) \subset \Omega.$$

The following Hadamard type theorem was proved by S. Granlund in [2].

2.7. Theorem. *Let u be a subminimum of*

$$I(u) = \int_{\Omega} F(x, \nabla u(x)) dm,$$

$r_1 < r < r_2$, and $\overline{B}^n(r_2) \subset \Omega$. Then u is bounded from above, and there is a constant

$$\lambda = \lambda(n, p, r/r_1, r_2/r, \alpha/\beta),$$

$0 < \lambda < 1$ such that

$$M(r) \leq \lambda M(r_1) + (1 - \lambda)M(r_2).$$

Since p -harmonic functions minimize (see e.g. [4, p. 59]) the integral

$$I(u) = \int_{\Omega} |\nabla u|^p dm,$$

Theorem 2.7 is related to Theorem 2.1 with $k = n$.

3. PRELIMINARIES

We start by recalling some basic properties of the Sobolev spaces from [4]. Let Ω be a nonempty open set in $\mathbb{R}^n$.

3.1. Lemma. [4, Theorem 1.24] *Let $u \in W_0^{1,p}(\Omega)$ and $v \in W^{1,p}(\Omega)$ be bounded. Then $uv \in W_0^{1,p}(\Omega)$.*

3.2. Lemma. [4, Lemma 3.11] *If $v \in W^{1,p}(\Omega)$ is a weak solution of (1.2) in Ω , then*

$$\int_{\Omega} \langle |\nabla v|^{p-2} \nabla v, \nabla \varphi \rangle dm = 0$$

for all $\varphi \in W_0^{1,p}(\Omega)$.

3.3. Theorem. [1, p. 99] *Let $f: \mathbb{R}^n \rightarrow \mathbb{R}$ be a locally Lipschitz mapping. Let $E \subset \mathbb{R}^n$ be an n -measurable set and $g: E \rightarrow \mathbb{R}$ be a nonnegative measurable function. Then*

$$(3.4) \quad \int_E g(x) |\nabla f(x)| dx_1 \cdots dx_n = \int_{\mathbb{R}} \left(\sum_{x \in f^{-1}(y)} g(x) \right) d\mathcal{H}^n(y).$$

3.5. Lemma. *Let $1 < p < \infty$, $0 < r < d_k(x)$ and fix an integer $1 \leq k \leq n$. Then*

$$u(x) = \int_r^{d_k(x)} s^{\frac{1-k}{p-1}} ds$$

is a solution of (1.2), i.e.

$$\sum_{i=1}^n \left\{ \frac{\partial}{\partial x_i} (u_{x_i} [u_{x_1}^2 + \cdots + u_{x_n}^2]^{\frac{p-2}{2}}) \right\} = 0.$$

Proof. We note that

$$\frac{\partial}{\partial x_i} d_k(x) = \frac{x_i}{d_k(x)},$$

and hence $u_{x_i} = x_i d_k(x)^{\frac{1-k}{p-1}-1}$. Then

$$\begin{aligned} u_{x_i} (u_{x_1}^2 + \dots + u_{x_k}^2)^{\frac{p-2}{2}} &= x_i d_k(x)^{\frac{1-k}{p-1}-1} \left[d_k(x)^{\frac{2(1-k)}{p-1}-2} \left(\sum_{j=1}^k x_j^2 \right) \right]^{\frac{p-2}{2}} \\ &= x_i d_k(x)^{\frac{1-k}{p-1}-1} d_k(x)^{\frac{(1-k)(p-2)}{p-1}} = x_i d_k(x)^{-k}. \end{aligned}$$

It follows that

$$\begin{aligned} \sum_{i=1}^k \frac{\partial}{\partial x_i} (x_i d_k(x)^{-k}) &= \sum_{i=1}^k d_k(x)^{-k} - k \sum_{i=1}^k x_i^2 d_k(x)^{-k-2} \\ &= k d_k(x)^{-k} - k d_k(x)^{-k-2} \left(\sum_{i=1}^k x_i^2 \right) = 0. \end{aligned}$$

□

Next we will prove two lemmas which are used later in the proof of Theorem 2.1.

3.6. Lemma. *Let $a > b > 0$, $p > 1$. Then*

$$(3.7) \quad C_1 \frac{a^{p-1} - b^{p-1}}{a - b} \leq \frac{a^{p-1} + b^{p-1}}{a + b} \leq C_2 \frac{a^{p-1} - b^{p-1}}{a - b},$$

with some constants $C_1, C_2 > 0$.

Proof. We examine the function

$$g_1(x) = \frac{(x^{p-1} + 1)(x - 1)}{(x^{p-1} - 1)(x + 1)}, \quad x > 1.$$

It is clear that

$$(3.8) \quad \lim_{x \rightarrow 1} g_1(x) = \frac{1}{p-1}, \quad \lim_{x \rightarrow \infty} g_1(x) = 1.$$

It is sufficient to find positive bounds for $g_1(x)$ for $x > 1$. We will prove that the bounds are in fact given by (3). First we note that

$$\begin{cases} (p-2)(x^p - 1) + p(x - x^{p-1}) < 0, & \text{for } p \in (1, 2), \\ (p-2)(x^p - 1) + p(x - x^{p-1}) = 0, & \text{for } p = 2, \\ (p-2)(x^p - 1) + p(x - x^{p-1}) > 0, & \text{for } p > 2, \end{cases}$$

and

$$\begin{cases} x - x^{p-1} < 0, & \text{for } p \in (1, 2), \\ x - x^{p-1} = 0, & \text{for } p = 2, \\ x - x^{p-1} > 0, & \text{for } p > 2. \end{cases}$$

Hence

$$\begin{cases} g_1(x) \in (1, 1/(p-1)), & \text{for } p \in (1, 2), \\ g_1(x) = 1, & \text{for } p = 2, \\ g_1(x) \in (1/(p-1), 1), & \text{for } p > 2. \end{cases}$$

□

3.9. Lemma. *Let $a > b > 0$. Then*

$$(3.10) \quad C_3(a^{p-2} + b^{p-2}) \leq \frac{a^{p-1} - b^{p-1}}{a - b} \leq C_4(a^{p-2} + b^{p-2}),$$

for $p \geq 2$, and

$$(3.11) \quad C_3(a^{2-p} + b^{2-p})^{-1} \leq \frac{a^{p-1} - b^{p-1}}{a - b} \leq C_4(a^{2-p} + b^{2-p})^{-1},$$

for $p \in (1, 2]$ with some constants $C_3, C_4 > 0$.

Proof. The proof is similar to that of Lemma 3.6. First we study the function

$$g_2(x) = \frac{x^{p-1} - 1}{(x - 1)(x^{p-2} + 1)}.$$

As in Lemma 3.6, it is sufficient for (3.10) to find positive bounds for $g_2(x)$ for $x > 0$. We note that $\lim_{x \rightarrow 1} g_2(x) = (p - 1)/2$ and $\lim_{x \rightarrow \infty} g_2(x) = 1$. We obtain

$$\begin{cases} (p - 3)(1 - x^{p-1}) + (p - 1)x(1 - x^{p-3}) < 0, & \text{for } p \in (1, 3), \\ (p - 3)(1 - x^{p-1}) + (p - 1)x(1 - x^{p-3}) = 0, & \text{for } p = 3, \\ (p - 3)(1 - x^{p-1}) + (p - 1)x(1 - x^{p-3}) > 0, & \text{for } p > 3, \end{cases}$$

and

$$\begin{cases} x(x^{p-3} - 1) < 0, & \text{for } p \in (1, 3), \\ x(x^{p-3} - 1) = 0, & \text{for } p = 3, \\ x(x^{p-3} - 1) > 0, & \text{for } p > 3. \end{cases}$$

It follows that

$$\begin{cases} g_2(x) \in ((p - 1)/2, 1), & \text{for } p \in (1, 3), \\ g_2(x) = 1, & \text{for } p = 3, \\ g_2(x) \in (1, (p - 1)/2), & \text{for } p > 3. \end{cases}$$

To prove (3.11) we study the function

$$g_3(x) = \frac{(x^{p-1} - 1)(x^{2-p} + 1)}{x - 1}.$$

Now $\lim_{x \rightarrow 1} g_3(x) = 2(p - 1)$ and $\lim_{x \rightarrow \infty} g_3(x) = 1$. Again, we have

$$\begin{cases} (-2p + 3)(x - 1) + (x^{p-1} - x^{2-p}) < 0, & \text{for } p \in (1, 3/2), \\ (-2p + 3)(x - 1) + (x^{p-1} - x^{2-p}) = 0, & \text{for } p = 3/2, \\ (-2p + 3)(x - 1) + (x^{p-1} - x^{2-p}) > 0, & \text{for } p > 3/2, \end{cases}$$

and

$$\begin{cases} x^{p-1} - x^{2-p} < 0, & \text{for } p \in (1, 3/2), \\ x^{p-1} - x^{2-p} = 0, & \text{for } p = 3/2, \\ x^{p-1} - x^{2-p} > 0, & \text{for } p > 3/2, \end{cases}$$

and thus

$$\begin{cases} g_3(x) \in (2(p - 1), 1), & \text{for } p \in (1, 3/2), \\ g_3(x) = 1, & \text{for } p = 3/2, \\ g_3(x) \in (1, 2(p - 1)), & \text{for } p > 3/2. \end{cases}$$

□

4. PROOF OF THEOREM 2.1

Suppose the contrary, that is, there exists $x_0 \in D_{r,R}$ such that

$$(4.1) \quad v(x_0) > (M(R) - M(r))u(x_0) + M(r),$$

or

$$v_{r,R}(x_0) > u(x_0).$$

Fix some $\varepsilon_0 > 0$, for which

$$v_{r,R}(x_0) - u(x_0) > \varepsilon_0.$$

Consider the set

$$U = \{x \in D_{r,R} : v_{r,R}(x) - u(x) > \varepsilon_0\} \neq \emptyset.$$

Choose a component O of U such that $x_0 \in O$. It is clear that $\overline{O} \cap \partial D_{r,R} = \emptyset$ and $(v_{r,R}(x) - u(x))|_{\partial O} = 0$. Fix $\varepsilon_2 > \varepsilon_1 > 0$ and the balls $O_1 = B_k(x_0, \varepsilon_1)$, $O_2 = B_k(x_0, \varepsilon_2)$. Let $\varphi(x) = \eta(d_k(x))$ be a locally Lipschitz function with the properties:

$$(4.2) \quad \begin{cases} \varphi \equiv 1 & \text{for all } x \in O_1, \\ \varphi \equiv 0 & \text{for all } x \in D_{r,R} \setminus O_2. \end{cases}$$

Then the function $\psi = (v_{r,R}(x) - u(x))\varphi^2$ has a support $\text{supp } \psi \subset \overline{O}_2$ and by Lemma 3.1 $\psi \in W_0^{1,p}(\Omega)$ for all Ω such that $\text{supp } \psi \subset \Omega$. Since $v_{r,R}$ and u are generalized solutions of (1.2) we have by Lemma 3.2

$$\begin{aligned} & \int_{\text{supp } \psi} \langle \nabla \psi, |\nabla v_{r,R}|^{p-2} \nabla v_{r,R} - |\nabla u|^{p-2} \nabla u \rangle dm \\ &= \int_{\text{supp } \psi} \langle \nabla \psi, |\nabla v_{r,R}|^{p-2} \nabla v_{r,R} \rangle dm - \int_{\text{supp } \psi} \langle \nabla \psi, |\nabla u|^{p-2} \nabla u \rangle dm = 0. \end{aligned}$$

Next, we note that

$$\nabla \psi = \varphi^2(\nabla v_{r,R} - \nabla u) + 2\varphi(v_{r,R} - u)\nabla \varphi.$$

Thus, we may write

$$\begin{aligned} 0 &= \int_{\text{supp } \psi} \langle \nabla \psi, |\nabla v_{r,R}|^{p-2} \nabla v_{r,R} - |\nabla u|^{p-2} \nabla u \rangle dm \\ &= \int_{O \cap O_2} \langle \nabla \psi, |\nabla v_{r,R}|^{p-2} \nabla v_{r,R} - |\nabla u|^{p-2} \nabla u \rangle dm \\ &= \int_{O \cap O_2} \varphi^2 \langle \nabla v_{r,R} - \nabla u, |\nabla v_{r,R}|^{p-2} \nabla v_{r,R} - |\nabla u|^{p-2} \nabla u \rangle dm \\ &\quad + 2 \int_{O \cap O_2} \varphi(v_{r,R} - u) \langle \nabla \varphi, |\nabla v_{r,R}|^{p-2} \nabla v_{r,R} - |\nabla u|^{p-2} \nabla u \rangle dm \end{aligned}$$

or

$$\begin{aligned} \int_{O \cap O_2} \varphi^2 \langle \nabla v_{r,R} - \nabla u, |\nabla v_{r,R}|^{p-2} \nabla v_{r,R} - |\nabla u|^{p-2} \nabla u \rangle dm \\ = -2 \int_{O \cap O_2} \varphi(v_{r,R} - u) \langle \nabla \varphi, |\nabla v_{r,R}|^{p-2} \nabla v_{r,R} - |\nabla u|^{p-2} \nabla u \rangle dm \end{aligned}$$

or

$$\begin{aligned} (4.3) \quad & \left| \int_{O \cap O_2} \varphi^2 \langle \nabla v_{r,R} - \nabla u, |\nabla v_{r,R}|^{p-2} \nabla v_{r,R} - |\nabla u|^{p-2} \nabla u \rangle dm \right| \\ & \leq 2 \int_{O \cap O_2} |\varphi| |v_{r,R} - u| |\nabla \varphi| | |\nabla v_{r,R}|^{p-2} \nabla v_{r,R} - |\nabla u|^{p-2} \nabla u | dm. \end{aligned}$$

Let

$$\Phi(\lambda) = |\nabla(\lambda v_{r,R} + (1 - \lambda)u)|^{p-2} \nabla(\lambda v_{r,R} + (1 - \lambda)u)$$

for $\lambda \in [0, 1]$, and note that

$$\Phi(0) = |\nabla u|^{p-2} \nabla u \quad \text{and} \quad \Phi(1) = |\nabla v_{r,R}|^{p-2} \nabla v_{r,R}.$$

Now we write

$$\begin{aligned} (4.4) \quad & |\nabla v_{r,R}|^{p-2} \nabla v_{r,R} - |\nabla u|^{p-2} \nabla u = \Phi(1) - \Phi(0) = \int_0^1 \Phi'(\lambda) d\lambda \\ & = \int_0^1 \left[(\nabla v_{r,R} - \nabla u) |\nabla(\lambda v_{r,R} + (1 - \lambda)u)|^{p-2} + (p - 2) \nabla(\lambda v_{r,R} + (1 - \lambda)u) \right. \\ & \quad \cdot |\nabla(\lambda v_{r,R} + (1 - \lambda)u)|^{p-4} \langle \nabla v_{r,R} - \nabla u, \nabla(\lambda v_{r,R} + (1 - \lambda)u) \rangle \left. \right] d\lambda, \end{aligned}$$

and obtain

$$\begin{aligned} (4.5) \quad & \langle \nabla v_{r,R} - \nabla u, |\nabla v_{r,R}|^{p-2} \nabla v_{r,R} - |\nabla u|^{p-2} \nabla u \rangle \\ & = |\nabla v_{r,R} - \nabla u|^2 \int_0^1 |\nabla(\lambda v_{r,R} + (1 - \lambda)u)|^{p-2} d\lambda \\ & \quad + (p - 2) \int_0^1 |\nabla(\lambda v_{r,R} + (1 - \lambda)u)|^{p-4} \langle \nabla v_{r,R} - \nabla u, \nabla(\lambda v_{r,R} + (1 - \lambda)u) \rangle^2 d\lambda. \end{aligned}$$

If $p \geq 2$ then

$$\begin{aligned} (4.6) \quad & \langle \nabla v_{r,R} - \nabla u, |\nabla v_{r,R}|^{p-2} \nabla v_{r,R} - |\nabla u|^{p-2} \nabla u \rangle \\ & \geq |\nabla v_{r,R} - \nabla u|^2 \int_0^1 |\nabla(\lambda v_{r,R} + (1 - \lambda)u)|^{p-2} d\lambda. \end{aligned}$$

If $p < 2$, we have

$$\begin{aligned} & |\nabla v_{r,R} - \nabla u|^2 \int_0^1 |\nabla(\lambda v_{r,R} + (1-\lambda)u)|^{p-2} d\lambda \\ & + (p-2) \int_0^1 |\nabla(\lambda v_{r,R} + (1-\lambda)u)|^{p-4} \langle \nabla v_{r,R} - \nabla u, \nabla(\lambda v_{r,R} + (1-\lambda)u) \rangle^2 d\lambda \\ & \geq (p-1) |\nabla v_{r,R} - \nabla u|^2 \int_0^1 |\nabla(\lambda v_{r,R} + (1-\lambda)u)|^{p-2} d\lambda. \end{aligned}$$

This together with (4.5) gives

$$\begin{aligned} (4.7) \quad & \langle \nabla v_{r,R} - \nabla u, |\nabla v_{r,R}|^{p-2} \nabla v_{r,R} - |\nabla u|^{p-2} \nabla u \rangle \\ & \geq (p-1) |\nabla v_{r,R} - \nabla u|^2 \int_0^1 |\nabla(\lambda v_{r,R} + (1-\lambda)u)|^{p-2} d\lambda, \quad 1 < p \leq 2. \end{aligned}$$

It follows from (4.4) that for every $p > 1$,

$$(4.8) \quad |\nabla v_{r,R}|^{p-2} \nabla v_{r,R} - |\nabla u|^{p-2} \nabla u \leq C_5 |\nabla v_{r,R} - \nabla u| \int_0^1 |\nabla(\lambda v_{r,R} + (1-\lambda)u)|^{p-2} d\lambda,$$

at every point where $v_{r,R}$ has differential. Here $C_5 = 1 + |p-2|$. Setting

$$I(p) = \int_0^1 |\nabla(\lambda v_{r,R} + (1-\lambda)u)|^{p-2} d\lambda$$

and using (4.3), (4.6), (4.7) and (4.8) we obtain

$$(4.9) \quad \int_{O \cap O_2} \varphi^2 I(p) |\nabla v_{r,R} - \nabla u|^2 dm \leq C_6 \int_{O \cap O_2} I(p) |\varphi| |v_{r,R} - u| |\nabla \varphi| |\nabla v_{r,R} - \nabla u| dm,$$

where $C_6 = 2C_5 / \min\{1, p-1\}$.

We note that

$$|\nabla(\lambda v_{r,R} + (1-\lambda)u)|^2 = \lambda^2 |\nabla v_{r,R}|^2 + 2\lambda(1-\lambda) \langle \nabla v_{r,R}, \nabla u \rangle + (1-\lambda)^2 |\nabla u|^2,$$

and therefore

$$(4.10) \quad |\lambda |\nabla v_{r,R}| - (1-\lambda) |\nabla u|| \leq |\nabla(\lambda v_{r,R} + (1-\lambda)u)| \leq \lambda |\nabla v_{r,R}| + (1-\lambda) |\nabla u|$$

for an arbitrary $\lambda \in [0, 1]$. Let $p \geq 2$. We suppose that $|\nabla v_{r,R}| > |\nabla u|$. Then by (4.10),

$$\begin{aligned} (4.11) \quad I(p) & \leq \int_0^1 (\lambda (|\nabla v_{r,R}| - |\nabla u|) + |\nabla u|)^{p-2} d\lambda \\ & = \frac{1}{|\nabla v_{r,R}| - |\nabla u|} \int_{|\nabla u|}^{|\nabla v_{r,R}|} s^{p-2} ds = \frac{1}{p-1} \frac{|\nabla v_{r,R}|^{p-1} - |\nabla u|^{p-1}}{|\nabla v_{r,R}| - |\nabla u|}. \end{aligned}$$

Next by (4.10),

$$\begin{aligned}
I(p) &\geq \int_0^1 |\lambda |\nabla v_{r,R}| - (1-\lambda) |\nabla u||^{p-2} d\lambda \\
&= \int_0^1 |\lambda (|\nabla v_{r,R}| + |\nabla u|) - |\nabla u||^{p-2} d\lambda \\
&= \int_s^1 (\lambda (|\nabla v_{r,R}| + |\nabla u|) - |\nabla u|)^{p-2} d\lambda \\
&\quad + \int_0^s (|\nabla u| - \lambda (|\nabla v_{r,R}| + |\nabla u|))^{p-2} d\lambda,
\end{aligned}$$

where

$$(4.12) \quad s = \frac{|\nabla u|}{|\nabla v_{r,R}| + |\nabla u|}.$$

By computing both of the last two integrals, we obtain

$$(4.13) \quad I(p) \geq \frac{1}{p-1} \frac{|\nabla v_{r,R}|^{p-1} + |\nabla u|^{p-1}}{|\nabla v_{r,R}| + |\nabla u|}.$$

Let $1 < p < 2$. As above, we assume $|\nabla v_{r,R}| > |\nabla u|$. Then by (4.10),

$$\begin{aligned}
I(p) &\leq \int_0^1 |\lambda |\nabla v_{r,R}| - (1-\lambda) |\nabla u||^{2-p} d\lambda \\
&= \int_0^1 |\lambda (|\nabla v_{r,R}| + |\nabla u|) - |\nabla u||^{2-p} d\lambda \\
&= \int_0^s (|\nabla u| - \lambda (|\nabla v_{r,R}| + |\nabla u|))^{2-p} d\lambda \\
&\quad + \int_s^1 (\lambda (|\nabla v_{r,R}| + |\nabla u|) - |\nabla u|)^{2-p} d\lambda,
\end{aligned}$$

where s is defined in (4.12), and hence

$$(4.14) \quad I(p) \leq \frac{1}{p-1} \frac{|\nabla v_{r,R}|^{p-1} + |\nabla u|^{p-1}}{|\nabla v_{r,R}| + |\nabla u|}.$$

By (4.11), it follows that

$$(4.15) \quad I(p) \geq \frac{1}{p-1} \frac{|\nabla v_{r,R}|^{p-1} - |\nabla u|^{p-1}}{|\nabla v_{r,R}| - |\nabla u|}.$$

Setting $a = |\nabla v_{r,R}|$ and $b = |\nabla u|$ in (3.7), (3.10) and (3.11), we can obtain by (4.11), (4.13), (4.14) and (4.15), for $p \geq 2$

$$(4.16) \quad C_7 (|\nabla v_{r,R}|^{p-2} + |\nabla u|^{p-2}) \leq I(p) \leq C_8 (|\nabla v_{r,R}|^{p-2} + |\nabla u|^{p-2}),$$

or

$$(4.17) \quad C_7 (|\nabla v_{r,R}|^{2-p} + |\nabla u|^{2-p})^{-1} \leq I(p) \leq C_8 (|\nabla v_{r,R}|^{2-p} + |\nabla u|^{2-p})^{-1},$$

$1 < p \leq 2$, with some constants $C_7, C_8 > 0$. The case $|\nabla v_{r,R}| < |\nabla u|$ is analogous. This may be written as

$$(4.18) \quad C_9 (|\nabla v_{r,R}|^{|p-2|} - |\nabla u|^{|p-2|}) \leq I(p) \leq C_{10} (|\nabla v_{r,R}|^{|p-2|} + |\nabla u|^{|p-2|}),$$

where $C_9 = \min\{C_7, 1/C_8\}$ and $C_{10} = \max\{1/C_7, C_8\}$.

Thus by (4.9), (4.18) we find,

$$(4.19) \quad \begin{aligned} & \int_{O \cap O_2} \varphi^2 |\nabla v_{r,R} - \nabla u|^2 (|\nabla v_{r,R}|^{|p-2|} + |\nabla u|^{|p-2|}) dm \\ & \leq C_{11} \int_{O \cap O_2} |\varphi| |v_{r,R} - u| |\nabla \varphi| |\nabla v_{r,R} - \nabla u| (|\nabla v_{r,R}|^{|p-2|} + |\nabla u|^{|p-2|}) dm \\ & \leq C_{11} \left(\int_{O \cap O_2} |\nabla \varphi|^2 |v_{r,R} - u|^2 (|\nabla v_{r,R}|^{|p-2|} + |\nabla u|^{|p-2|}) dm \right)^{1/2} \\ & \quad \cdot \left(\int_{O \cap O_2} \varphi^2 |\nabla v_{r,R} - \nabla u|^2 (|\nabla v_{r,R}|^{|p-2|} + |\nabla u|^{|p-2|}) dm \right)^{1/2} \end{aligned}$$

and

$$\begin{aligned} & \int_{O \cap O_2} \varphi^2 |\nabla v_{r,R} - \nabla u|^2 (|\nabla v_{r,R}|^{|p-2|} + |\nabla u|^{|p-2|}) dm \\ & \leq C_{11}^2 \int_{O \cap O_2} |\nabla \varphi|^2 |v_{r,R} - u|^2 (|\nabla v_{r,R}|^{|p-2|} + |\nabla u|^{|p-2|}) dm. \end{aligned}$$

Remembering (4.2) we have

$$\begin{aligned} & \int_{O \cap O_1} |\nabla v_{r,R} - \nabla u|^2 (|\nabla v_{r,R}|^{|p-2|} + |\nabla u|^{|p-2|}) dm \\ & \leq C_{11}^2 \int_{D_{r,R} \cap (O_2 \setminus \overline{O_1})} |\nabla \varphi|^2 |v_{r,R} - u|^2 (|\nabla v_{r,R}|^{|p-2|} + |\nabla u|^{|p-2|}) dm. \end{aligned}$$

Because φ is constant on $\Sigma_k(t)$ and $|\nabla d_k| \equiv 1$, we have by Theorem 3.3

$$\begin{aligned} & \int_{D_{r,R} \cap (O_2 \setminus \overline{O_1})} |\nabla \varphi|^2 |v_{r,R} - u|^2 (|\nabla v_{r,R}|^{|p-2|} + |\nabla u|^{|p-2|}) dm \\ & \leq \int_{\{x: \varepsilon_1 < d_k(x) < \varepsilon_2\}} |\nabla \varphi|^2 |v_{r,R} - u|^2 (|\nabla v_{r,R}|^{|p-2|} + |\nabla u|^{|p-2|}) dm \\ & = \int_{\varepsilon_1}^{\varepsilon_2} \eta'^2 H(t) dt, \end{aligned}$$

where

$$(4.20) \quad H(t) = \int_{\Sigma_k(t)} |v_{r,R} - u|^2 (|\nabla v_{r,R}|^{p-2} + |\nabla u|^{p-2}) d\mathcal{H}^{n-1}.$$

By Hölder's inequality

$$1 \leq \int_{\varepsilon_1}^{\varepsilon_2} \eta' H(t)^{1/2} H(t)^{-1/2} dt \leq \left(\int_{\varepsilon_1}^{\varepsilon_2} \eta'^2 H(t) dt \right)^{1/2} \left(\int_{\varepsilon_1}^{\varepsilon_2} H^{-1}(t) dt \right)^{1/2}.$$

It follows that

$$(4.21) \quad \left(\int_{\varepsilon_1}^{\varepsilon_2} H^{-1}(t) dt \right)^{-1} \leq \int_{\varepsilon_1}^{\varepsilon_2} \eta'^2 H(t) dt,$$

for all $\varphi(x) = \eta(d_k(x))$ satisfying (4.2).

We define a function $\hat{\eta}$ by the formula

$$\hat{\eta}(s) = \left(\int_{\varepsilon_1}^s H^{-1}(t) dt \right) \left(\int_{\varepsilon_1}^{\varepsilon_2} H^{-1}(t) dt \right)^{-1}.$$

Now $\hat{\eta}(\varepsilon_1) = 0$ and $\hat{\eta}(\varepsilon_2) = 1$. Because

$$\hat{\eta}'(s) = \frac{1}{H(s)} \left(\int_{\varepsilon_1}^{\varepsilon_2} \frac{dt}{H(t)} \right)^{-1},$$

we have by (4.21)

$$\begin{aligned} \left(\int_{\varepsilon_1}^{\varepsilon_2} H^{-1}(t) dt \right)^{-1} &\leq \inf_{\varphi} \int_{\varepsilon_1}^{\varepsilon_2} \eta'^2 H(t) dt \\ &\leq \int_{\varepsilon_1}^{\varepsilon_2} \hat{\eta}'^2 H(t) dt = \left(\int_{\varepsilon_1}^{\varepsilon_2} H^{-1}(t) dt \right)^{-1}. \end{aligned}$$

Because

$$\begin{aligned} &\int_{\varepsilon_1}^{\varepsilon_2} H^{-1}(t) dt \\ &= \int_{\varepsilon_1}^{\varepsilon_2} dt \left(\int_{\Sigma_k(t)} |v_{r,R} - u|^2 (|\nabla v_{r,R}|^{p-2} + |\nabla u|^{p-2}) d\mathcal{H}^{n-1} \right)^{-1} \rightarrow \infty, \end{aligned}$$

as $\varepsilon_2 \rightarrow \infty$, the claim follows. $\square$

5. PROOFS OF THE COROLLARIES

Proof of Corollary 2.4. Let

$$(5.1) \quad H(t) = \int_{\Sigma_k(t)} |v_{r,R} - u|^2 (|\nabla v_{r,R}|^{p-2} + |\nabla u|^{p-2}) d\mathcal{H}^{n-1}.$$

By Hölder's inequality

$$(S - r)^2 = \left(\int_r^S dt \right)^2 = \left(\int_r^S \frac{H^{-1/2}(t)}{H^{-1/2}(t)} dt \right)^2 \leq \left(\int_r^S H^{-1}(t) dt \right) \left(\int_r^S H(t) dt \right).$$

Hence

$$(5.2) \quad (S-r)^2 \left(\int_r^S H^{-1}(t) dt \right)^{-1} \leq \left(\int_r^S H(t) dt \right).$$

Now by (5.2) and Theorem 3.3

$$\begin{aligned} & \left[\int_r^S dt \left(\int_{\Sigma_k(t)} |v_{r,R} - u|^2 (|\nabla v_{r,R}|^{p-2} + |\nabla u|^{p-2}) d\mathcal{H}^{n-1} \right)^{-1} \right]^{-1} \\ & \leq \frac{1}{(S-r)^2} \int_{D_{r,S}} |v_{r,R} - u|^2 (|\nabla v_{r,R}|^{p-2} + |\nabla u|^{p-2}) dm \rightarrow 0, \end{aligned}$$

as $S \rightarrow \infty$, proving the claim. $\square$

Proof of Corollary 2.6. Since

$$\int_{D_{r,\infty}} |v_{r,R} - u|^2 (|\nabla v_{r,R}|^{p-2} + |\nabla u|^{p-2}) dm \leq M < \infty,$$

we have for $S > r$,

$$\frac{1}{S^2} \int_{D_{r,S}} |v_{r,R} - u|^2 (|\nabla v_{r,R}|^{p-2} + |\nabla u|^{p-2}) dm \leq \frac{M}{S^2} \rightarrow 0,$$

as $S \rightarrow \infty$. $\square$

REFERENCES

- [1] L.C. EVANS AND R.F. GARIEPY: *Measure theory and fine properties of functions*. Studies in Advanced Mathematics, CRC Press, Boca Raton – New York – London – Tokyo, 1992.
- [2] S. GRANLUND: *Three-Circles Theorem For Variational Integrals*. Universität Bonn Preprint no. 507, April 1982.
- [3] J. HADAMARD: *Sur les fonctions entières*. C.R. Acad. Sci. Paris 122 (1896), 1257–1258.
- [4] J. HEINONEN, T. KILPELÄINEN AND O. MARTIO: *Nonlinear potential theory of degenerate elliptic equations*. Clarendon Press 1993.
- [5] V. MAZ'YA AND T. SHAPOSHNIKOVA: *Jacques Hadamard, a universal mathematician*. History of Mathematics, 14, American Mathematical Society, Providence, RI; London Mathematical Society, London, 1998.
- [6] V.M. MIKLYUKOV AND M. VUORINEN: *A generalized maximum principle for the differences of p -harmonic functions*. Proceedings on Analysis and Geometry (Russian) (Novosibirsk Akademgorodok, 1999), 401–413, Izdat. Ross. Akad. Nauk Sib. Otd. Inst. Mat., ISBN 5-86134-091-9, Novosibirsk, 2000.
- [7] M. PROTTER AND H. WEINBERGER: *Maximum Principles in Differential Equations*, Springer-Verlag, New York, 1984.
- [8] R.M. ROBINSON: *Hadamard's three circles theorem*, Bull. Amer. Math. Soc. 50 (1944), 795–802.

(Vladimir M. Miklyukov) DEPARTMENT OF MATHEMATICS, VOLGOGRAD STATE UNIVERSITY, UNIVERSITETSKII PROSPECT 100, VOLGOGRAD 400062, RUSSIA, FAX + TEL: +7-8442 471608
E-mail address: miklyuk@mail.ru

(Antti Rasila) DEPARTMENT OF MATHEMATICS AND SYSTEMS ANALYSIS, AALTO UNIVERSITY, P.O. BOX 1110 0, FI-00076 AALTO, FINLAND, FAX +358-9-451 3016
E-mail address: antti.rasila@iki.fi

(Matti Vuorinen) DEPARTMENT OF MATHEMATICS, FI-20014 UNIVERSITY OF TURKU, FINLAND

E-mail address: `vuorinen@utu.fi`